

Ferroelectric phase transition, ionicity condensation, and multicriticality in charge transfer organic complexes

Jun-ichiro Kishine,^{a*} Tadeusz Luty,^b and Kenji Yonemitsu,^a

^a *Department of Theoretical Studies,
Institute for Molecular Science, Okazaki 444-8585, Japan*
and

*Department of Functional Molecular Science,
Graduate University for Advanced Studies,
Okazaki 444-8585, Japan*

^b *Institute of Physical and Theoretical Chemistry Technical University of Wrocław 50-370 Wrocław, Poland*

(Dated: February 2, 2008)

To elucidate a novel pressure-temperature phase diagram of the quasi-one-dimensional mixed-stack charge-transfer (CT) complex TTF-CA, we study the quasi-one-dimensional spin-1 Blume-Emery-Griffith (BEG) model. In addition to the local charge transfer energy and the inter-stack polar (dipole-dipole) interaction, we take account of the inter-stack electrostriction effect. Using the self-consistent chain-mean-field theory, where the intra-stack degrees of freedom are exactly treated by the transfer-matrix method, we reproduce the gas-liquid-solid like phase diagram corresponding to the neutral (N), paraelectric ionic (I_{para}), and ferroelectric ionic (I_{ferro}) phases, respectively. We also give an explanation on the experimentally observed multicritical behavior and concomitant discontinuous inter-stack lattice contraction in TTF-CA.

PACS numbers:

I. INTRODUCTOIN

In the “critical phase control technology,” condensed molecular materials play quite a promising role, because molecular orbitals and stacking architecture are manipulable in a desirable way. To elucidate interrelation of constituent molecular structures and emergence of various thermodynamic phases such as superconductivity, magnetism, and ferroelectricity is of great interest there. A neutral-to-ionic phase transition (NIT) in quasi-one-dimensional charge-transfer (CT) complexes comprising mixed-stack architecture of electron donor (D) and acceptor (A) molecules¹ has played a key role in this field.

In particular, phase control by pressure^{2,3} or laser radiation^{3,4} in the tetrathiafulvalene-*p*-chloranil (TTF-CA), that exhibits the NIT around 80K at ambient pressure, has attracted a great deal of interest. Very recently, Collet *et al.*,⁵ using highly refined time-resolved X-ray diffraction technique, have reported direct observation of a photo-induced paraelectric-to-ferroelectric structural order in the crystal. In the ionic phase, the D^+A^- pair forms a dimer due to the electrostatic instability⁶ or subsequent spin-Peierls instability.⁷ The ionized dimer on the DA chain carries a local electric dipole moment p with opposite directions depending on the dimerization patterns $\underline{D^+A^-}$ or $\underline{A^-D^+}$. Once p acquires a macroscopic mean value $\eta = \langle p \rangle \neq 0$, a spontaneous inversion symmetry breaking (SISB) occurs and the system undergoes a phase transition to a ferroelectric-ionic (I_{ferro}) phase.

The ionic phase itself is simply described by ionicity condensation $c = \langle p^2 \rangle \sim 1$. Since η is a symmetry-breaking order parameter but c is not, we expect that η and c play separate roles. The appearance of two distinct order parameters η and c is a direct consequence of the degeneracy of the two configurations of dimerization pattern, $IA (\cdots \underline{D^+A^-} \underline{D^+A^-} \cdots)$ and $(\cdots \underline{A^-D^+} \underline{A^-D^+} \cdots)$.

Recently, the respective roles of c and η have been highlighted in both equilibrium² and nonequilibrium³ processes. Using the neutron diffraction along with nuclear-quadrupole-resonance (NQR) measurements, Lemée-Cailleau *et al.*² found a novel phase where *the system is ionic but dipoles remain disordered*, i.e., a paraelectric ionic (I_{para}) phase. They proposed a pressure-temperature phase diagram of TTF-CA, where the N, I_{para} , and I_{ferro} phases are like gas, liquid, and solid phases, respectively. The ferroelectric order is well signaled by the appearance of $(0, 2k+1, 0)$ Bragg peaks that indicate the inversion symmetry breaking. The “sublimation” line separating the N and I_{ferro} phases continues up to a triple point $(P_t, T_t) \sim (500\text{MPa}, 210\text{K})$. Above the triple point, in addition to the “crystallization (or melting)” line, there appears a “condensation” line separating the I_{para} and I_{ferro} phases accompanied by a concomitant discontinuous change of c , ending at a critical point $(P_c, T_c) \sim (700\text{MPa}, 250\text{K})$. The purpose of this paper is to give a qualitative understanding of this phase diagram. This phase diagram, in addition to the phenomenological description offered in ref.³.

Since the SISB is prohibited by thermal fluctuations in a purely one-dimensional stack, inter-stack coupling is required to realize the SISB. In addition, the experimental observation strongly indicates that electronic and lattice degrees of freedom are coupled with each other in a unique manner. That is to say, upon crossing the tran-

*Address after September 2003: *Department of Physics, Faculty of Engineering, Kyushu Institute of Technology, 1-1 Sensuicho, Tobata, Kitakyushu 804-8550, Japan*

sition lines in the gas-liquid-solid like phase diagram,^{2,3} the unit cell parameter b (for the axis perpendicular to the stack) exhibits about 0.5 % discontinuous contraction at the condensation transition but exhibits continuous contraction at the crystallization transition. On the other hand, the unit cell parameter a (for the stacking axis) exhibits only continuous contraction at the condensation transition and below it remains almost constant.³ Now we are ready to ask the question: (1) what kind of inter-stack interactions are responsible for the occurrence of the I_{ferro} phase, and (2) how the lattice anomalies are coupled to the phase transitions?

As for the first question, Luty *et al.*^{3,8} stressed that the inter-stack non-polar coupling⁷ alone cannot drive the ferroelectric ordering and the *dipolar* coupling plays an essential role. As for the second question, Kawamoto *et al.*⁹ took account of the charge distribution on the atoms inside each molecule by an ab initio quantum chemical method and elucidated the importance of inter-stack Coulomb attraction $\sim -0.14\text{eV}$, which may cause inter-stack electrostriction (Coulomb-lattice coupling).

II. QUASI-ONE-DIMENSIONAL BLUME-EMERY-GRIFFITH MODEL AND INTERCHAIN MEAN FIELD THEORY

Now we shall set up a model. The ground-state energy of the mixed stacks has three minima as a function of the dimerization displacement, i.e., the N and the degenerate IA and IB states. The three states may be described by the spin-1 Ising variable $p_{i,j} = 0, \pm 1$ on the i th dimer inside the j th stack.^{2,3,8} The charge transfer energy (Δ), the intra- (with subscript $\parallel$) and inter- (with subscript $\perp$) stack dipolar (J) and non-polar (K) interactions, and the coupling with the electric field (E) are described by the quasi one-dimensional (Q1D) Blume-Emery-Griffith (BEG) model,¹⁰ $\mathcal{H} = \mathcal{H}_{\parallel} + \mathcal{H}_{\perp}$, where

$$\mathcal{H}_{\parallel} = - \sum_{i,j} [J_{\parallel} p_{i,j} p_{i+1,j} + K_{\parallel} p_{i,j}^2 p_{i+1,j}^2 - \Delta p_{i,j}^2 - E p_{i,j}], \quad (1)$$

$$\mathcal{H}_{\perp} = - \sum_{i,j} [J_{\perp} p_{i,j} p_{i,j+1} + K_{\perp} p_{i,j}^2 p_{i,j+1}^2]. \quad (2)$$

The intra-stack dipolar interaction $J_{\parallel}$ is caused by coupling between the charge transfer and the lattice distortion,⁸ while the inter-stack dipolar interaction is regarded as a direct interaction between the induced dipoles on adjacent stacks. The intra-stack couplings are much stronger than the inter-stack couplings, and the electric dipoles are aligned along the stacks. The energy cost to create one D^+A^- pair is given in the limit of no molecular overlap by $\Delta = I - A - \alpha V$, where I and A denote the donor's ionization energy and the acceptor's affinity, respectively, and αV denotes the Madelung energy.¹¹ Generally speaking, increasing pressure decreases the lattice spacing a and consequently in-

creases V . Therefore, Δ decreases upon applying pressure.

We treat the Hamiltonian (2) by using the self-consistent chain mean-field theory¹². Introducing the thermal averages, $\eta = \langle p_{i,j} \rangle$ and $c = \langle p_{i,j}^2 \rangle$, we have the effective 1D BEG model,

$$\mathcal{H}_{\parallel}^{\text{eff}} = - \sum_i [J_{\parallel} p_i p_{i+1} + K_{\parallel} p_i^2 p_{i+1}^2 - \tilde{\Delta} p_i^2 - \tilde{E} p_i] + \frac{z_{\perp}}{2} N J_{\perp} \eta^2 + \frac{z_{\perp}}{2} N K_{\perp} c^2, \quad (3)$$

where $\tilde{\Delta} = \Delta - z_{\perp} K_{\perp} c$ and $\tilde{E} = E - z_{\perp} J_{\perp} \eta$, with $z_{\perp} = 2$ being the inter-stack coordination number. Treating $\mathcal{H}_{\parallel}^{\text{eff}}$ exactly by the transfer matrix method, we obtain the free energy per site,

$$f_{\text{BEG}}(\eta, c, T) = -T \ln \lambda(\tilde{\Delta}, \tilde{E}, T) + J_{\perp} \eta^2 + K_{\perp} c^2, \quad (4)$$

where $\lambda(\tilde{\Delta}, \tilde{E}, T)$ is the maximum eigenvalue of the transfer matrix for $\mathcal{H}_{\parallel}^{\text{eff}}$, given by

$$\mathbf{T} = \begin{pmatrix} e^{\beta(J_{\parallel} + K_{\parallel} - \tilde{\Delta} - \tilde{E})} & e^{-\beta(\tilde{\Delta} + \tilde{E})/2} & e^{\beta(-J_{\parallel} + K_{\parallel} - \tilde{\Delta})} \\ e^{-\beta(\tilde{\Delta} + \tilde{E})/2} & 1 & e^{-\beta(\tilde{\Delta} - \tilde{E})/2} \\ e^{\beta(-J_{\parallel} + K_{\parallel} - \tilde{\Delta})} & e^{-\beta(\tilde{\Delta} - \tilde{E})/2} & e^{\beta(J_{\parallel} + K_{\parallel} - \tilde{\Delta} + \tilde{E})} \end{pmatrix}, \quad (5)$$

with $\beta = 1/T$. Possible phase diagrams of the BEG model have been extensively studied through mean-field theories,^{10,13} renormalization-group,¹⁴ and transfer-matrix methods.¹⁵ For the parameter regions relevant to the present case, $J_{\parallel}$, $J_{\perp}$, $K_{\parallel}$, $K_{\perp}$, and Δ are all positive, so that a solid-liquid-gas type phase diagram with proper slopes of transition lines is not obtained.

III. INTERCHAIN ELECTROSTRICTION

Then, we consider the inter-stack lattice degrees of freedom that have not explicitly been taken into account in (2). It is well known that an electrostriction effect potentially converts a continuous transition to a discontinuous one, since this gives rise to an additional negative free-energy term that contains the forth power of the relevant order parameter.¹⁶ In the present case, we phenomenologically introduce an additional free energy,

$$f_{\text{elst}}(c, y) = -\frac{c^2}{b_0 + y} + \frac{1}{2} k y^2, \quad (6)$$

where the first and second terms represent Coulomb attraction between the nearest neighbor stacks⁹ and the elastic energy for the distortion in the inter-stack direction. Note that $f_{\text{elst}}(c, 0)$ has already been absorbed into $K_{\perp}$. The lattice constant without distortion is b_0 , and y denotes the distortion. By minimizing $f_{\text{elst}}(c, y)$ with respect to y , we obtain the optimized lattice constant,

$$b(T) = b_0 + y(T) \sim b_0 - 2\varepsilon_{\text{elst}} b_0^2 c^2, \quad (7)$$

where $\varepsilon_{\text{elst}} = 1/(2kb_0^4)$ is a small constant. We thus have the energy gain due to the lattice distortion,

$$f_{\text{elst}}(c, T) \sim -\varepsilon_{\text{elst}} c^4. \quad (8)$$

Now, solving the self-consistent equations is reduced to searching (c, η) that gives the absolute minimum of the total free energy, $f(\eta, c, T) = f_{\text{BEG}}(\eta, c, T) + f_{\text{elst}}(c, T)$. The ionic phase is characterized by the ionicity condensation $c = 1$, while the ferroelectric phase is characterized by $\eta \neq 0$.

Note that, in the present scheme, any phase with $c \neq 1$ is regarded as “neutral” and that the neutral phase has always $\eta = 0$. In the BEG model, because of three states $p_{i,j} = 0, \pm 1$, c approaches the universal constant $c = 2/3$ in the high temperature limit, where the entropy term dominates the internal energy term. Therefore, in the parameter region where c continuously increases upon decreasing temperature, we have $2/3 < c \leq 1$. In the experiments,¹ the ionicity continuously increases upon decreasing temperature and jumps from $c \sim 0.3$ to $c \sim 0.6$ at the NIT. Thus, concerning the quantitative magnitude of the ionicity, there arises a difference between the experimental result and the present analysis. This apparent difference comes from the fact that we mapped the intra-stack CT transfer and the DA dimerization onto the simple spin-1 Ising variables. Therefore, we should regard the difference as an artifact of the classical BEG model.

IV. NUMERICAL RESULTS AND PHASE DIAGRAM

From now on, we set $J_{\parallel} = 1$ as an energy unit and the electric field E is set to be zero. In Fig. 1, we show the temperature dependence of c and η for various magnitudes of Δ with $K_{\parallel} = 0.4$, $K_{\perp} = 0.06$, $J_{\perp} = 0.03$, and $\varepsilon_{\text{elst}} = 0.0095$. We introduce the condensation temperature T_{cond} and the crystallization temperature T_{cryst} . The ionicity jumps into $c = 1$ at T_{cond} , while the ferroelectric order parameter acquires a finite magnitude $\eta \neq 0$ at T_{cryst} . The ground state becomes ionic for $\Delta < 1.49$. Both c and η exhibits a discontinuous change at the same transition temperature (the sublimation temperature) for $1.42 < \Delta < 1.49$. That is to say, $T_{\text{cond}} = T_{\text{cryst}}$. For $\Delta < 1.42$, there appears a region, $T_{\text{cryst}} < T < T_{\text{cond}}$, where the system is ionic but still paraelectric. This region is identified with the I_{para} phase that is observed in TTF-CA under pressure. The point ($\Delta_t = 1.42, T_t = 0.45$) is identified with the *triple point* (indicated by “TP” in Fig. 5). For $\Delta < 1.42$, c still exhibits a discontinuous change at T_{cond} , but η continuously evolves at T_{cryst} , as shown in Fig. 1(c). The discontinuity jump of c becomes smaller and seems to vanish, as Δ decreases, as shown in Fig. 1(d). We stress that this discontinuity is a direct consequence of the weak but finite electrostriction effect. Without the electrostriction, as Δ decreases, T_{cond} and T_{cryst} continue to coincide with each

other, and the transition simply changes from discontinuous to continuous at some critical value of Δ .^{10,13,14,15}




FIG. 1: Temperature dependence of c and η for various magnitudes of Δ with $K_{\parallel} = 0.4$, $K_{\perp} = 0.06$, $J_{\perp} = 0.03$, and $\varepsilon_{\text{elst}} = 0.0095$. The condensation and crystallization temperature, T_{cond} and T_{cryst} , respectively, are indicated. Locations of the Δ values in (a)-(d) are indicated in the phase diagram of Fig. 5.

As clearly seen from (7), about 0.5% discontinuous contraction of the inter-stack lattice constant (unit cell parameter b) is accompanied by the discontinuous jump of the ionicity. In Fig. 2, setting b_0 as a length unit, we show the temperature dependence of the unit cell parameter b given by (7), using the same parameter set as that in Fig. 1. Although the magnitude of the discontinuous contraction depends on the parameter choice of $\varepsilon_{\text{elst}}$ and b_0 , the qualitative nature (b jumps at T_{cond}) does not change.

To see the discontinuity of the ionicity more closely, we show in Fig. 3 the Δ dependence of the discontinuity at the condensation temperature, Δc . It is clearly seen that Δc decreases as Δ decreases and eventually reaches zero at $\Delta = 1.25$. For $\Delta < 1.25$, the condensation occurs without ionicity jump. Then, the lattice contraction at T_{cond} also becomes continuous. Therefore, $\Delta = 1.25$ with the corresponding $T_{\text{cond}} = 0.76$ is identified with a *critical point* (indicated by “CP” in Fig. 5). This result is well consistent with the experimental fact that the ionicity jump finishes at the critical point.²

The dielectric constant is given by $\varepsilon = 1 + 4\pi\alpha$, where the uniform polarizability is $\alpha = \frac{1}{T} \sum_{i,j} \sum_{l,m} [\langle p_{i,j} p_{l,m} \rangle - \langle p_{i,j} \rangle \langle p_{l,m} \rangle] = (c - \eta^2)/T$. In Fig. 4, we show the temperature dependence of α for various magnitudes of Δ . It is seen that along the N- I_{ferro} boundary, the polarizability exhibits a sharp single cusp at $T_{\text{cond}} = T_{\text{cryst}}$. For $\Delta_c < \Delta < \Delta_t$, a discontinuous jump occurs at $T = T_{\text{cond}}$ and a cusp at $T = T_{\text{cryst}}$. The discontinuity at $T = T_{\text{cond}}$ finishes at $\Delta = \Delta_c$.




FIG. 2: Temperature dependence of the unit cell parameter b , $b(T)$. Locations of the Δ values in (a)-(d) are indicated in the phase diagram of Fig. 5.




FIG. 3: Temperature dependence of the discontinuity of the ionicity, Δc .




FIG. 4: Temperature dependence of the polarizability, α . Locations of the Δ values in (a)-(d) are indicated in the phase diagram of Fig. 5.

In Fig. 5, we show the phase diagram of the system for $K_{\parallel} = 0.4$, $K_{\perp} = 0.06$, $J_{\perp} = 0.03$, and $\varepsilon_{\text{elst}} = 0.0095$. Regarding the decreasing Δ as increasing pressure, this phase diagram is consistent with the experimentally found, pressure-temperature phase diagram of TTF-CA.² The triple point, the critical point, and the observed inter-stack lattice contraction are reproduced. For simplicity, we here ignored the change of Δ due to thermal lattice contraction. Exactly speaking, to convert our Δ - T phase diagram to a P - T diagram, we need to take account of the temperature dependence of Δ , $\Delta(T)$. By appropriately treating $\Delta(T)$, we may obtain the corresponding P - T phase diagram satisfying the Clausius-Clapeyron relation. We stress that, even when we take this simple view, a qualitative nature of the phase diagram is not changed. Identifying the triple point $(\Delta_t, T_t) = (1.42, 0.45)$ with the experimentally obtained one $(P_t, T_t) \sim (500\text{MPa}, 210\text{K})$, we see that our parameter choice here corresponds to $K_{\perp} = 28\text{K}$ and $J_{\perp} = 14\text{K}$.

Lajzerowicz and Sivardi re¹⁷ extensively developed a mean-field analysis of the BEG model and obtained liquid-gas-solid like phase diagrams on the P - T plane. However, they considered a lattice gas analogue of a simple fluid, where the physical pressure of the lattice gas is simply given by $-f$, with f being the Helmholtz free energy per volume. In the present context, the pressure of the spin system has no physical meaning and the phase diagram obtained by Lajzerowicz and Sivardi re cannot be applied to TTF-CA.

V. CONCLUDING REMARKS

In this paper, we showed that the inter-stack electrostriction causes the $-c^4$ term with a small coefficient and makes the phase diagram rich. We thus conclude that the inter-stack polar interaction together with the inter-stack electrostriction drives the discontinuous inter-stack lattice contraction and the multicritical behavior observed in TTF-CA.^{2,3} Our considerations are limited to classical models, i.e., we considered classical *effective* models, where all the microscopic (electronic or phonon) degrees of freedom are implicitly integrated out. To obtain the truly microscopic mechanism behind the multicriticality, we need to get back to such a microscopic Hamiltonian as an extended Peierls-Hubbard model.¹⁸ We would keep this issue for future study.

This work was partly supported by a Grant-in-Aid for Scientific Research (C) from Japan Society for the Promotion of Science. TL wishes to acknowledge hospitality at the Institute for Molecular Sciences, Okazaki, and thank colleagues for creating a pleasant and stimulating environment there.




FIG. 5: Phase diagram for $K_{\parallel} = 0.4$, $K_{\perp} = 0.06$, $J_{\perp} = 0.03$, and $\epsilon_{\text{elst}} = 0.0095$. The solid and dashed lines represent discontinuous and continuous transitions, respectively. **TP** and **CP** represent the triple point and the critical point, respectively. Locations of the Δ values used in (a)-(d) of Figs. 1, 2, and 4 are indicated by the horizontal arrows.

-
- ¹ J. B. Torrance, J. E. Vazquez, J. J. Mayerle, and V. Y. Lee, Phys. Rev. Lett. **46**, 253 (1981).
 - ² M. H. Lemée-Cailleau, M. Le Cointe, H. Cailleau, T. Luty, F. Moussa, J. Roos, D. Brinkmann, B. Toudic, C. Ayache, and N. Karl, Phys. Rev. Lett. **79**, 1690 (1997).
 - ³ T. Luty, H. Cailleau, S. Koshihara, E. Collet, M. Takesada, M-H. Lemée-Cailleau, M. Buron-Le Cointe, N. Nagaosa, Y. Tokura, E. Zienkiewicz, and B. Ouladdiaf, Europhys. Lett. **59**, 619 (2002).
 - ⁴ S. Koshihara, Y. Tokura, K. Takeda, T. Koda, Phys. Rev. Lett. **68**, 1148 (1992).
 - ⁵ E. Collet, M. H. Lemée-Cailleau, M. Buron-Le Cointe, H. Cailleau, M. Wulff, T. Luty, S. Koshihara, M. Meyer, L. Toupet, P. Rabiller and S. Techert, Science, **300**, 612 (2003).
 - ⁶ T. Iizuka-Sakano and Y. Toyozawa, J. Phys. Soc. Jpn. **65**, 671 (1996).
 - ⁷ N. Nagaosa, J. Phys. Soc. Jpn. **55**, 3488 (1986).
 - ⁸ T. Luty in *Relaxations of Excited States and Photo-Induced Structural Phase Transitions*, edited by K. Nasu, Springer Series in Solid-State Sciences **124** (Springer-Verlag, 1996) p. 142.
 - ⁹ T. Kawamoto, T. Iizuka-Sakano, Y. Shimoi, and S. Abe, Phys. Rev. B **64**, 205107 (2001).
 - ¹⁰ M. Blume, V. J. Emery, and R. B. Griffiths, Phys. Rev. A **4**, 1071 (1971).
 - ¹¹ H. M. McConnell, B. M. Hoffman, and R. M. Metzger, Proc. Natl. Acad. Sci. U.S.A. **53**, 46 (1965).
 - ¹² D. J. Scalapino, Y. Imry, and P. Pincus, Phys. Rev. B **11**, 2042 (1975).
 - ¹³ W. Hoston, A. N. Berker, Phys. Rev. Lett. **67**, 1027 (1991).
 - ¹⁴ A. N. Berker and M. Wortis, Phys. Rev. B **14**, 4946 (1976).
 - ¹⁵ E. Albayrak and M. Keskin, J. Mag. Mag. Mater. **213**, 201 (2000).
 - ¹⁶ See, for example, P. W. Anderson, *Basic Notions of Condensed Matter Physics*, (Addison-Wesley, 1984), Section 2 C.
 - ¹⁷ J. Lajzerowicz and J. Sivardière, Phys. Rev. A **11**, 2079 (1975).
 - ¹⁸ N. Miyashita, M. Kuwabara, and K. Yonemitsu, cond-mat/0211487 (unpublished), J. Phys. Soc. Jpn. (in press).